# Orthopaedic Sports Specialists, P.C.

Michael E. Joyce, M.D., Alex G. Dukas, M.D, Christian Merrill M.D/MBA 84 Glastonbury Blvd., Suite 101, Glastonbury, Connecticut 06033 Phone: 860-652-8883, Fax: 860-652-8887

## **Post Operative Rehabilitation:**

<u>Elbow Arthroscopy</u> (Posterior Compartment/Valgus Extension Overload, Plica)

#### Phase I: Immediate Motion Phase

Goals: Improve/regain full range of motion Decrease pain/inflammation Retard muscular atrophy

#### A. Day 1-4

Range of motion to tolerance (Extension/Flexion and Supination/Pronation) \*Often full elbow extension is not capable due to pain. <u>Gentle</u> overpressure into extension Wrist Flex/ext stretches Gripping exercises (putty) Isometrics wrist ext/flex Isometrics elbow ext/flex Compression dressing , ice 4-5 times daily

## B. Day 5-10

Range of motion to tolerance (at least 20-90 degrees) Overpressure into extension Joint mobilization to re-establish ROM-Wrist flex/ext stretches Continue isometrics Continue use of ice, compression to control swelling

#### C. Day 11-14

Range of motion exercises to tolerance

(at least 10-100 degrees) Overpressure into extension (3-4 times daily) Continue joint mobilization techniques initiate light dumbbell program (PRE's) Biceps, triceps, wrist flex/ext, sup/pronators Continue use of ice post-exercise

## Phase II Intermediate Phase

## <u>Goals</u>: Improve strength/power/endurance Increase range of motion Initiate functional activities

## A. Week 2-4

Full range of motion exercises (4-5 times daily) Overpressure into elbow extension Continue PRE program for elbow & wrist musculature Initiate shoulder program (esp ER,RTC) Continue joint mobilization Continue ice post-exercise

#### B. Week 4-7

Continue all exercises listed above Initiate <u>light</u> upper body program Continue use of ice post-activity

## Phase III Advanced Strengthening Program

<u>Goals</u>: Improve strength/power/endurance Gradual Return to functional activities

## **Criteria to enter Phase III:**

- 1) Full non-painful range of motion
- 2) Strength 75% > of contralateral side
- 3) No pain or tenderness

## A. Week 8-12

Continue PRE program for elbow & wrist Continue shoulder program Continue stretching or elbow/shoulder Initiate interval program & gradually Return to sport activities.